arXiv:math/9904017v4 [math.QA] 18 Jun 1999

Drinfeld Basis And a Nonclassical Free Boson Representation of Twisted Quantum Affine Superalgebra $U_q[osp(2|2)^{(2)}]$

Wen-Li Yang [1] and Yao-Zhong Zhang [2]

[1] Institute of Modern Physics, Northwest University, Xian 710069, China

[2] Department of Mathematics, University of Queensland, Brisbane, Qld 4072, Australia

Abstract

We derive from the super RS algebra the Drinfeld basis of the twisted quantum affine superalgebra $U_q[osp(2|2)^{(2)}]$ by means of the Gauss decomposition technique. We explicitly construct a nonclassical level-one representation of $U_q[osp(2|2)^{(2)}]$ in terms of two q-deformed free boson fields.

Keywords: Drinfeld basis, quantum affine superalgebras, free boson representation.

1 Introduction

The algebraic analysis method based on infinite dimensional non-abelian symmetries has proved extremely successful in both formulating and solving lower dimensional integrable systems [1][2]. The key elements of this method are infinite-dimensional highest weight representations of quantum (super) algebras and vertex operators. Drinfeld bases of quantum (super) algebras are of great importance in constructing the infinite-dimensional representations and vertex operators.The Drinfeld bases of quantum affine bosonic algebras were given by Drinfeld [3]. For the case of quantum affine superalgebras, the Drinfeld bases have been known only for $U_q[gl(m|n)^{(1)}]$ [4, 5, 6] and $U_q[osp(1|2)^{(1)}]$ [7, 8]. In this paper, we derive the Drinfeld basis of the twisted quantum affine superalgebra $U_q[osp(2|2)^{(2)}]$

by using the super version of the Reshetikhin-Semenov-Tian-Shansky (RS) algebra [9] the Gauss decomposition techinque of Ding-Frenkel [10]. Moreover, we explicitly construct a nonclassical level-one representation of $U_q[osp(2|2)^{(2)}]$ by means of two q-deformed free boson fields.

2 Drinfeld Basis of $U_q[osp(2|2)^{(2)}]$

The symmetric Cartan matrix of the twisted affine superalgebra $osp(2|2)^{(2)}$ is (a_{ij}) with $a_{11} = a_{22} = 1,\ a_{12} = a_{21} = -1$. Twisted quantum affine superalgebra $U_q[osp(2|2)^{(2)}]$ is a q-analogue of the universal enveloping algebra of $osp(2|2)^{(2)}$ and is generated by the Chevalley generators $\{E_i,\ F_i,\ K_i^{\pm 1}|i = 0, 1\}$. The $\mathbf{Z}_2$-grading of the Chevalley generators is $[E_i] = [F_i] = 1,\ [K_i] = 0,\ i = 0, 1$. The defining relations are

$$\begin{aligned}
&K_iK_j = K_jK_i, \\
&K_jE_iK_j^{-1} = q^{a_{ij}}E_i, \qquad K_jF_iK_j^{-1} = q^{-a_{ij}}F_i, \\
&\{E_i, F_j\} = \delta_{ij}\frac{K_i - K_i^{-1}}{q - q^{-1}}
\end{aligned} \tag{2.1}$$

plus q-Serre relations which we omit. Here $\{X, Y\} \equiv XY + YX$.

$U_q[osp(2|2)^{(2)}]$ is a $\mathbf{Z}_2$-graded quasi-triangular Hopf algebra endowed with the coproduct Δ, counit ϵ and antipode S given by

$$\begin{aligned}
&\Delta(E_i) = E_i \otimes 1 + K_i \otimes E_i, \qquad \Delta(F_i) = F_i \otimes K_i^{-1} + 1 \otimes F_i, \\
&\Delta(K_i) = K_i \otimes K_i, \qquad \epsilon(E_i) = \epsilon(F_i) = 0, \qquad \epsilon(K_i) = 1, \\
&S(E_i) = -K_i^{-1}E_i, \qquad S(F_i) = -F_iK_i, \qquad S(K_i) = K_i^{-1}, \quad i = 0, 1.
\end{aligned} \tag{2.2}$$

The antipode S is a $\mathbf{Z}_2$-graded algebra anti-homomorphism, i.e. for homogeneous elements $a, b \in U_q[osp(2|2)^{(2)}]$, we have $S(ab) = (-1)^{[a][b]}S(b)S(a)$.

We now give our definition of $U_q[osp(2|2)^{(2)}]$ in terms of Drinfeld generators.

Definition 1 *$U_q[osp(2|2)^{(2)}]$ is an associative algebra with unit 1 and the Drinfeld generators: $X^\pm(z)$ and $\psi^\pm(z)$, a central element c and a nonzero complex parameter q. $\psi^\pm(z)$ are invertible. The gradings of the generators are: $[X^\pm(z)] = 1$ and $[\psi^\pm(z)] = [c] = 0$. The relations are given by*

$$\begin{aligned}
&\psi^\pm(z)\psi^\pm(w) = \psi^\pm(w)\psi^\pm(z), \\
&\psi^+(z)\psi^-(w) = \frac{(z_+q + w_-)(z_- + w_+q)}{(z_+ + w_-q)(z_-q + w_+)}\psi^-(w)\psi^+(z), \\
&\psi^\pm(z)^{-1}X^+(w)\psi^\pm(z) = \frac{z_\pm + wq}{z_\pm q + w}X^+(w),
\end{aligned}$$

$$
\begin{aligned}
&\psi^{\pm}(z)X^{-}(w)\psi^{\pm}(z)^{-1} = \frac{z_{\mp} + wq}{z_{\mp}q + w}X^{-}(w),\\
&(z + wq^{\pm 1})X^{\pm}(z)X^{\pm}(w) + (zq^{\pm 1} + w)X^{\pm}(w)X^{\pm}(z) = 0,\\
&\{X^{+}(z), X^{-}(w)\} = \frac{1}{(q - q^{-1})zw}\left[\delta(\frac{w}{z}q^{c})\psi^{+}(w_{+}) - \delta(\frac{w}{z}q^{-c})\psi^{-}(z_{+})\right].
\end{aligned}
\tag{2.3}
$$

Expand the currents in the form

$$
\begin{aligned}
X^{\pm}(z) &= \sum_{n\in \mathbf{Z}} X^{\pm}_{n}\, z^{-n-1},\\
\psi^{\pm}(z) &= \sum_{n\in \mathbf{Z}} \psi^{\pm}_{n}\, z^{-n} = K^{\pm 1}\exp\left(\pm(q - q^{-1})\sum_{n>0} H_{\pm n}z^{\mp n}\right).
\end{aligned}
\tag{2.4}
$$

In terms of modes $\{H_n \,|\, n \in \mathbf{Z} - \{0\},\ X^{\pm}_{n} \,|\, n \in \mathbf{Z}\}$ and K, the defining relations of $U_q[osp(2|2)^{(2)}]$ are given by

$$
\begin{aligned}
&c: \ \text{central element},\\
&[K, H_n] = 0, \qquad KX^{\pm}_{n}K^{-1} = q^{\pm 1}X^{\pm}_{n},\\
&[H_n, H_m] = \delta_{n+m,0}(-1)^{n}\frac{[n]_q[nc]_q}{n}, \quad n \neq 0,\\
&[H_n, X^{\pm}_{m}] = \pm\frac{1}{n}(-1)^{n}[n]_q q^{\mp\frac{c}{2}|n|}X^{\pm}_{n+m}, \quad n \neq 0,\\
&X^{\pm}_{n+1}X^{\pm}_{m} + q^{\pm 1}\,X^{\pm}_{m}X^{\pm}_{n+1} + q^{\pm 1}\,X^{\pm}_{n}X^{\pm}_{m+1} + X^{\pm}_{m+1}X^{\pm}_{n} = 0,\\
&\{X^{+}_{n}, X^{-}_{m}\} = \frac{1}{q - q^{-1}}\left(q^{\frac{c}{2}(n-m)}\psi^{+}_{n+m} - q^{\frac{c}{2}(m-n)}\psi^{-}_{n+m}\right).
\end{aligned}
\tag{2.5}
$$

Here and throughout $[i]_q = (q^{i} - q^{-i})/(q - q^{-1})$.

The Chevalley generators are obtained by the formulae:

$$
\begin{aligned}
&K_1 = K, \qquad E_1 = X^{+}_{0}, \qquad F_1 = X^{-}_{0},\\
&K_0 = q^{c}K^{-1}, \qquad E_0 = X^{-}_{1}K^{-1}, \qquad F_0 = -KX^{+}_{-1}.
\end{aligned}
\tag{2.6}
$$

In terms of the Chevalley generators, the Drinfeld generators can be built up recursively by

$$
\begin{aligned}
&H_1 = q^{\frac{c}{2}}K^{-1}(X^{+}_{0}X^{-}_{1} + X^{-}_{1}X^{+}_{0}),\\
&H_{-1} = q^{-\frac{c}{2}}K(X^{+}_{-1}X^{-}_{0} + X^{-}_{0}X^{+}_{-1}),\\
&X^{\pm}_{n+1} = \mp q^{\pm\frac{c}{2}}[H_1, X^{\pm}_{n}], \qquad X^{\pm}_{-n-1} = \mp q^{\pm\frac{c}{2}}[H_{-1}, X^{\pm}_{-n}], \quad n \geq 0
\end{aligned}
\tag{2.7}
$$

plus the formulae for $H_n,\ H_{-n}$ $(n > 0)$ given as follows

$$
H_{\pm n} = \pm\frac{1}{q - q^{-1}}\sum_{p_1+2p_2+\cdots+np_n=n}\frac{(-1)^{\sum p_i - 1}\,(\sum p_i - 1)!}{p_1!\cdots p_n!}(K^{\mp 1}\psi^{\pm}_{\pm 1})^{p_1}\cdots(K^{\mp 1}\psi^{\pm}_{\pm n})^{p_n},
\tag{2.8}
$$

where

$$
\psi^{\pm}_{\pm n} = \pm(q - q^{-1})q^{\mp\frac{c}{2}(n-2)}\{X^{\pm}_{\pm n\mp 1}, X^{\mp}_{\pm 1}\}, \quad n > 0.
\tag{2.9}
$$

3 Derivation of Drinfeld Basis from Super RS Algebra

Let us recall the definition of the super RS algebra. Let $R(z) \in End(V \otimes V)$, where V is a $\mathbf{Z}_2$ graded vector space, be a R-matrix which satisfies the graded Yang-Baxter equation

$$R_{12}(z)R_{13}(zw)R_{23}(w) = R_{23}(w)R_{13}(zw)R_{12}(z). \tag{3.10}$$

We introduce [5, 7]

Definition 2 *: The super RS algebra* $U(\mathcal{R})$ *is generated by invertible L-operators* $L^{\pm}(z)$*, which obey the relations*

$$\begin{aligned} R(\frac{z}{w})L_1^{\pm}(z)L_2^{\pm}(w) &= L_2^{\pm}(w)L_1^{\pm}(z)R(\frac{z}{w}), \\ R(\frac{z_+}{w_-})L_1^{+}(z)L_2^{-}(w) &= L_2^{-}(w)L_1^{+}(z)R(\frac{z_-}{w_+}), \end{aligned} \tag{3.11}$$

where $L_1^{\pm}(z) = L^{\pm}(z) \otimes 1$*,* $L_2^{\pm}(z) = 1 \otimes L^{\pm}(z)$ *and* $z_{\pm} = zq^{\pm\frac{c}{2}}$*. For the first formula of (3.11), the expansion direction of* $R(\frac{z}{w})$ *can be chosen in* $\frac{z}{w}$ *or* $\frac{w}{z}$*, but for the second formula, the expansion direction must only be in* $\frac{z}{w}$*.*

The multiplication rule for the tensor product is defined by

$$(a \otimes b)(a' \otimes b') = (-1)^{[b][a']}(aa' \otimes bb'), \tag{3.12}$$

for homogeneous elements $a,\ b,\ a',\ b'$ of $U_q[osp(2|2)^{(2)}]$.

In the following we apply the super RS algebra to derive the Drinfeld basis of $U_q[osp(2|2)^{(2)}]$. We take $R(\frac{z}{w})$ to be the R-matrix associated to the 3-dimensional representation V of $U_q[osp(2|2)^{(2)}]$. Let $v_1,\ v_2,\ v_3$ be the basis vectors of V with the $\mathbf{Z}_2$-grading $[v_1] = [v_3] = 0$ and $[v_2] = 1$. It can be shown that the R-matrix has the following form:

$$R(\frac{z}{w}) = \begin{pmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & a & 0 & b & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & d & 0 & c & 0 & r & 0 & 0 \\ 0 & f & 0 & a & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & g & 0 & e & 0 & c & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & a & 0 & b & 0 \\ 0 & 0 & s & 0 & g & 0 & d & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & f & 0 & a & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \end{pmatrix}, \tag{3.13}$$

where

$$\begin{aligned}
&a=\frac{q(z-w)}{zq^2-w}, \quad b=\frac{w(q^2-1)}{zq^2-w}, \quad c=-\frac{q^{1/2}w(q^2-1)(z-w)}{(zq^2-w)(zq+w)},\\
&d=\frac{q(z-w)(z+qw)}{(zq^2-w)(zq+w)}, \quad e=a-\frac{zw(q^2-1)(q+1)}{(zq^2-w)(zq+w)},\\
&f=\frac{z(q^2-1)}{zq^2-w}, \quad g=-\frac{q^{1/2}z(q^2-1)(z-w)}{(zq^2-w)(zq+w)},\\
&r=\frac{w^2(q-1)(q+1)^2}{(zq^2-w)(zq+w)}, \quad s=\frac{z^2(q-1)(q+1)^2}{(zq^2-w)(zq+w)}.
\end{aligned} \tag{3.14}$$

As in the non-super case [10], $L^{\pm}(z)$ allow the unique Gauss decomposition

$$L^{\pm}(z)=\left(\begin{array}{ccc}1&0&0\\ e_1^{\pm}(z)&1&0\\ e_{3,1}^{\pm}(z)&e_2^{\pm}(z)&1\end{array}\right)\left(\begin{array}{ccc}k_1^{\pm}(z)&0&0\\ 0&k_2^{\pm}(z)&0\\ 0&0&k_3^{\pm}(z)\end{array}\right)\left(\begin{array}{ccc}1&f_1^{\pm}(z)&f_{1,3}^{\pm}(z)\\ 0&1&f_2^{\pm}(z)\\ 0&0&1\end{array}\right), \tag{3.15}$$

where $e_{i,j}^{\pm}(z),\ f_{j,i}^{\pm}(z)$ and $k_i^{\pm}(z)$ $(i>j)$ are elements in the super RS algebra and $k_i^{\pm}(z)$ are invertible; and $e_i^{\pm}(z)\equiv e_{i+1,i}^{\pm}(z),\ f_i^{\pm}(z)\equiv f_{i,i+1}^{\pm}(z)$. Let us define

$$\begin{aligned}
X_i^{+}(z) &= f_i^{+}(z_+)-f_i^{-}(z_-),\\
X_i^{-}(z) &= e_i^{-}(z_+)-e_i^{+}(z_-).
\end{aligned} \tag{3.16}$$

By the definition of the super RS algebra and the Gauss decomposition formula (3.15), and after tedious calculations parallel to those of the $U_q[osp(1|2)^{(1)}]$ case, we arrive at

$$\begin{aligned}
k_i^{\pm}(z)k_j^{\pm}(w) &= k_j^{\pm}(w)k_i^{\pm}(z), \quad i,j=1,2,3,\\
k_1^{+}(z)k_1^{-}(w) &= k_1^{-}(w)k_1^{+}(z),\\
k_3^{+}(z)k_3^{-}(w) &= k_3^{-}(w)k_3^{+}(z),\\
\frac{z_\pm-w_\mp}{z_\pm q^2-w_\mp}k_1^{\pm}(z)k_2^{\mp}(w) &= \frac{z_\mp-w_\pm}{z_\mp q^2-w_\pm}k_2^{\mp}(w)k_1^{\pm}(z),\\
\frac{(z_\mp-w_\pm)(z_\mp+w_\pm q)}{(z_\mp q^2-w_\pm)(z_\mp q+w_\pm)}k_1^{\pm}(z)k_3^{\mp}(w)^{-1} &= \frac{(z_\pm-w_\mp)(z_\pm+w_\mp q)}{(z_\pm q^2-w_\mp)(z_\pm q+w_\mp)}k_3^{\mp}(w)^{-1}k_1^{\pm}(z),\\
\frac{(z_\pm-w_\mp q^2)(z_\pm q+w_\mp)}{(z_\pm q^2-w_\mp)(z_\pm+w_\mp q)}k_2^{\pm}(z)k_2^{\mp}(w) &= \frac{(z_\mp-w_\pm q^2)(z_\mp q+w_\pm)}{(z_\mp q^2-w_\pm)(z_\mp+w_\pm q)}k_2^{\mp}(w)k_2^{\pm}(z),\\
\frac{z_\pm-w_\mp}{z_\pm q^2-w_\mp}k_2^{\pm}(z)^{-1}k_3^{\mp}(w)^{-1} &= \frac{z_\mp-w_\pm}{z_\mp q^2-w_\pm}k_3^{\mp}(w)^{-1}k_2^{\pm}(z)^{-1},
\end{aligned}$$

$$\begin{aligned}
k_1^{\pm}(z)X_1^{-}(w)k_1^{\pm}(z)^{-1} &= \frac{z_\pm q^2-w}{q(z_\pm-w)}X_1^{-}(w),\\
k_1^{\pm}(z)^{-1}X_1^{+}(w)k_1^{\pm}(z) &= \frac{z_\mp q^2-w}{q(z_\mp-w)}X_1^{+}(w),
\end{aligned}$$

$$
\begin{aligned}
k_2^{\pm}(z)X_1^-(w)k_2^{\pm}(z)^{-1} &= \frac{(z_{\pm}q^2-w)(z_{\pm}+wq)}{q(z_{\pm}-w)(z_{\pm}q+w)}X_1^-(w),\\
k_2^{\pm}(z)^{-1}X_1^+(w)k_2^{\pm}(z) &= \frac{(z_{\mp}q^2-w)(z_{\mp}+wq)}{q(z_{\mp}-w)(z_{\mp}q+w)}X_1^+(w),\\
k_3^{\pm}(z)X_1^-(w)k_3^{\pm}(z)^{-1} &= \frac{z_{\pm}+wq}{z_{\pm}q+w}X_1^-(w),\\
k_3^{\pm}(z)^{-1}X_1^+(w)k_3^{\pm}(z) &= \frac{z_{\mp}+wq}{z_{\mp}q+w}X_1^+(w),\\
k_1^{\pm}(z)X_2^-(w)k_1^{\pm}(z)^{-1} &= \frac{z_{\pm}q+w}{z_{\pm}+wq}X_2^-(w),\\
k_1^{\pm}(z)^{-1}X_2^+(w)k_1^{\pm}(z) &= \frac{z_{\mp}q+w}{z_{\mp}+wq}X_2^+(w),\\
k_2^{\pm}(z)X_2^-(w)k_2^{\pm}(z)^{-1} &= \frac{(z_{\pm}-wq^2)(z_{\pm}q+w)}{q(z_{\pm}-w)(z_{\pm}+wq)}X_2^-(w),\\
k_2^{\pm}(z)^{-1}X_2^+(w)k_2^{\pm}(z) &= \frac{(z_{\mp}-wq^2)(z_{\mp}q+w)}{q(z_{\mp}-w)(z_{\mp}+wq)}X_2^+(w),\\
k_3^{\pm}(z)X_2^-(w)k_3^{\pm}(z)^{-1} &= \frac{z_{\pm}-wq^2}{q(z_{\pm}-w)}X_2^-(w),\\
k_3^{\pm}(z)^{-1}X_2^+(w)k_3^{\pm}(z) &= \frac{z_{\mp}-wq^2}{q(z_{\mp}-w)}X_2^+(w),
\end{aligned}
$$

$$
\begin{aligned}
(z-wq^2)X_1^+(z)X_2^+(w)+q(z-w)X_2^+(w)X_1^+(z) &= 0,\\
q(z-w)X_1^-(z)X_2^-(w)+(z-wq^2)X_2^-(w)X_1^-(z) &= 0,\\
(z+wq^{\pm 1})X_1^{\pm}(z)X_1^{\pm}(w)+(zq^{\pm 1}+w)X_1^{\pm}(w)X_1^{\pm}(z) &= 0,\\
(z+wq^{\pm 1})X_2^{\pm}(z)X_2^{\pm}(w)+(zq^{\pm 1}+w)X_2^{\pm}(w)X_2^{\pm}(z) &= 0,
\end{aligned}
$$

$$
\begin{aligned}
\{X_1^-(w),X_1^+(z)\} &= (q-q^{-1})\Big[-\delta(\frac{z}{w}q^{c})k_2^+(z_+)k_1^+(z_+)^{-1}\\
&\qquad +\delta(\frac{z}{w}q^{-c})k_2^-(w_+)k_1^-(w_+)^{-1}\Big],\\
\{X_2^-(w),X_2^+(z)\} &= (q-q^{-1})\Big[\delta(\frac{z}{w}q^{c})k_3^+(z_+)k_2^+(z_+)^{-1}\\
&\qquad -\delta(\frac{z}{w}q^{-c})k_3^-(w_+)k_2^-(w_+)^{-1}\Big],\\
\{X_2^-(w),X_1^+(z)\} &= (q-q^{-1})q^{-\frac{1}{2}}\Big[-\delta(-\frac{z}{w}q^{c-1})k_2^+(z_+)k_1^+(z_+)^{-1}\\
&\qquad +\delta(-\frac{z}{w}q^{-c-1})k_3^-(w_+)k_2^-(w_+)^{-1}\Big],\\
\{X_1^-(w),X_2^+(z)\} &= (q-q^{-1})q^{\frac{1}{2}}\Big[\delta(-\frac{z}{w}q^{c+1})k_2^+(-z_+q)k_1^+(-z_+q)^{-1}\\
&\qquad -\delta(-\frac{z}{w}q^{-c+1})k_3^-(-w_+q^{-1})k_2^-(-w_+q^{-1})^{-1}\Big],
\end{aligned}
\tag{3.17}
$$

where

$$
\delta(z)=\sum_{l\in\mathbf{Z}} z^l \tag{3.18}
$$

is a formal delta function which enjoys the following properties:

$$\delta(\frac{z}{w}) = \delta(\frac{w}{z}), \qquad \delta(\frac{z}{w})f(z) = \delta(\frac{z}{w})f(w). \tag{3.19}$$

Defining the algebraic homomorphism

$$\begin{aligned} X^{\pm}(z) &= z(q-q^{-1})\left[X_1^{\pm}(z) + X_2^{\pm}(-zq^{-1})\right], \\ \psi^-(z) &= (1+q^{-\frac{1}{2}}-q^{\frac{1}{2}})\phi_1(z) - \phi_2(-zq^{-1}), \\ \psi^+(z) &= \psi_1(z) - (1+q^{\frac{1}{2}}-q^{-\frac{1}{2}})\psi_2(-zq^{-1}), \end{aligned} \tag{3.20}$$

where $\phi_i(z) = k_{i+1}^+(z)k_i^+(z)^{-1}$, $\psi_i(z) = k_{i+1}^-(z)k_i^-(z)^{-1}$, $i = 1, 2$, then we obtain from (3.17), the current commutation relations (2.3).

4 Level Zero Representation

We consider the evaluation representation V_x of $U_q[osp(2|2)^{(2)}]$, where V is the 3-dimensional graded vector space with basis vectors $v_1,\ v_2$ and v_3. Let e_{ij} be the 3×3 matrix satisfying $e_{ij}v_k = \delta_{jk}v_i$. In the homogeneous gradation, the Chevalley generators of $U_q[osp(2|2)^{(2)}]$ are represented on V_x by

$$\begin{aligned} E_1 &= e_{12} - e_{23}, \quad F_1 = e_{21} + e_{32}, \quad K_1 = q^{e_{11}-e_{33}}, \\ E_0 &= xq^{-1}(-e_{21} + e_{32}), \quad F_0 = x^{-1}q(e_{12} + e_{23}), \quad K_0 = q^{-e_{11}+e_{33}}. \end{aligned} \tag{4.21}$$

Let V_x^{*S} denote the dual module of V_x, defined by $\pi_{V^*}(a) = \pi_V(S(a))^{st}$. On V_x^{*S}, the Chevalley generators are given by

$$\begin{aligned} E_1 &= -(q^{-1}e_{21} + e_{32}), \quad F_1 = qe_{12} - e_{23}, \quad K_1 = q^{-e_{11}+e_{33}}, \\ E_0 &= -xq^{-1}(e_{12} + q^{-1}e_{23}), \quad F_0 = x^{-1}q(-e_{21} + qe_{32}), \quad K_0 = q^{e_{11}-e_{33}}. \end{aligned} \tag{4.22}$$

The following proposition can be proved by induction.

Proposition 1 *: The Drinfeld generators are represented on V_x by*

$$\begin{aligned} H_m &= -x^m\frac{[m]_q}{m}\left[-(-1)^m q^{-m}e_{11} + (1-(-1)^m q^m)\, e_{22} + e_{33}\right], \\ X_m^+ &= -x^m\left(-(-1)^m e_{12} + q^{-m}e_{23}\right), \\ X_m^- &= x^m\left((-1)^m e_{21} + q^{-m}e_{32}\right), \qquad K = q^{e_{11}-e_{33}}, \end{aligned} \tag{4.23}$$

and on V_x^{*S} by

$$\begin{aligned} H_m &= -(-1)^m x^m\frac{[m]_q}{m}q^{-m}\left[e_{11} + \left(1-(-1)^m q^{-m}\right) e_{22} - (-1)^m q^{-m}e_{33}\right], \\ X_m^+ &= -(-1)^m x^m q^{-m}\left(q^{-m-1}e_{21} + (-1)^m e_{32}\right), \\ X_m^- &= (-1)^m x^m q^{-m}\left(q^{-m+1}e_{12} - (-1)^m e_{23}\right), \qquad K = q^{-e_{11}+e_{33}}. \end{aligned} \tag{4.24}$$

5 Nonclassical Free Boson Realization at Level One

We use the notation similar to that of [11, 12, 13]. Let us introduce two sets of bosonic oscillators $\{a_n,\ c_n,\ Q_a,\ Q_c | n \in \mathbf{Z}\}$ which satisfy the commutation relations

$$[a_n, a_m] = \delta_{n+m,0}(-1)^n \frac{[n]_q^2}{n}, \qquad [a_0, Q_a] = 1,$$
$$[c_n, c_m] = \delta_{n+m,0} \frac{[n]_q^2}{n}, \qquad [c_0, Q_c] = 1. \tag{5.25}$$

The remaining commutation relations are zero. Introduce the q-deformed free boson fields

$$a(z;\kappa) = Q_a + a_0 \ln z - \sum_{n\neq 0} \frac{a_n}{[n]_q} q^{\kappa|n|} z^{-n},$$
$$c(z) = Q_c + c_0 \ln z - \sum_{n\neq 0} \frac{c_n}{[n]_q} z^{-n} \tag{5.26}$$

and set

$$a_\pm(z) = \pm(q - q^{-1}) \sum_{n>0} a_{\pm n} z^{\mp n} \pm a_0 \ln q. \tag{5.27}$$

Then

Theorem 1 *: The Drinfeld generators of $U_q[osp(2|2)^{(2)}]$ at level one have the following nonclassical realization by the free boson fields*

$$\psi^\pm(z) = e^{a_\pm(z)},$$
$$X^\pm(z) =: e^{\pm a(z;\mp\frac{1}{2})}\, Y^\pm(z) : F^\pm, \tag{5.28}$$

where $F^+ = q^{1/2} + q^{-1/2}$, $F^- = 1/(q - q^{-1})$ and

$$Y^+(z) = e^{c(q^{1/2}z)} + e^{-c(-q^{-1/2}z)},$$
$$Y^-(z) = e^{c(-q^{1/2}z)} + e^{-c(q^{-1/2}z)}. \tag{5.29}$$

This free boson realization is nonclassical in the sense that $X^-(z)$ defined by (5.28) does not have classical or $q \to 1$ limit.

Proof. We prove this theorem by checking that the bosonized currents (5.28) satisfy the defining relations (2.3) of $U_q[osp(2|2)^{(2)}]$ with $c = 1$. It is easily seen that the first two relations in (2.3) are true by construction. The third and fourth ones follow from the definition of $X^\pm(z)$ and the commutativity between a_n and c_n. So we only need to check the last two relations in (2.3).

We write

$$Z^\pm(z) =: e^{\pm a(z;\mp\frac{1}{2})} : . \tag{5.30}$$

We obtain the operator products

$$\begin{aligned}
Z^{\pm}(z)Z^{\pm}(w) &= (z+q^{\mp 1}w)\ :Z^{\pm}(z)Z^{\pm}(w):, \\
Z^{+}(z)Z^{-}(w) &= (z+w)^{-1}\ :Z^{+}(z)Z^{-}(w):. \\
Y^{\pm}(z)Y^{\pm}(w) &= \pm q^{1/2}(z-w) : e^{c(\pm q^{1/2}z)}e^{c(\pm q^{1/2}w)} : \mp q^{-1/2}(z-w) : e^{-c(\mp q^{-1/2}z)}e^{-c(\mp q^{-1/2}w)} : \\
&\quad \pm\frac{q^{-1/2}}{z+q^{-1}w} : e^{c(\pm q^{1/2}z)}e^{-c(\mp q^{-1/2}w)} : \mp\frac{q^{1/2}}{z+qw} : e^{-c(\mp q^{-1/2}z)}e^{c(\pm q^{1/2}w)} :, \\
Y^{+}(z)Y^{-}(w) &= q^{1/2}(z+w) : e^{c(q^{1/2}z)}e^{c(-q^{1/2}w)} : -q^{-1/2}(z+w) : e^{-c(-q^{-1/2}z)}e^{-c(q^{-1/2}w)} : \\
&\quad +\frac{q^{-1/2}}{z-q^{-1}w} : e^{c(q^{1/2}z)}e^{-c(q^{-1/2}w)} : -\frac{q^{1/2}}{z-qw} : e^{-c(-q^{-1/2}z)}e^{c(-q^{1/2}w)} : .
\end{aligned}$$

Then the second last relation in (2.3) is easily seen to be true, and as to the last relation we have

$$\begin{aligned}
\{X^{+}(z),X^{-}(w)\} &= \frac{q^{1/2}+q^{-1/2}}{q-q^{-1}} : Z^{+}(z)Z^{-}(w) : \\
&\quad \left[\left(\frac{q^{-1/2}}{(z+w)(z-q^{-1}w)}+\frac{q^{1/2}}{(w+z)(w-qz)}\right) : e^{c(q^{1/2}z)}e^{-c(q^{-1/2}w)} : \right. \\
&\quad \left. -\left(\frac{q^{1/2}}{(z+w)(z-qw)}+\frac{q^{-1/2}}{(w+z)(w-q^{-1}z)}\right) : e^{-c(-q^{-1/2}z)}e^{c(-q^{1/2}w)} : \right] \\
&= \frac{1}{(q-q^{-1})zw} : Z^{+}(z)Z^{-}(w) : \\
&\quad \left[\left(\delta(-\frac{w}{z})-\delta(\frac{w}{z}q^{-1})\right) : e^{c(q^{1/2}z)}e^{-c(q^{-1/2}w)} : \right. \\
&\quad \left. -\left(\delta(-\frac{w}{z})-\delta(\frac{w}{z}q)\right) : e^{-c(-q^{-1/2}z)}e^{c(-q^{1/2}w)} : \right] \\
&= \frac{1}{(q-q^{-1})zw}\left(\delta(\frac{w}{z}q)-\delta(\frac{w}{z}q^{-1})\right) : Z^{+}(z)Z^{-}(w) : \\
&= \frac{1}{(q-q^{-1})zw}\left[\delta(\frac{w}{z}q)\psi^{+}(wq^{1/2})-\delta(\frac{w}{z}q^{-1})\psi^{-}(wq^{-1/2})\right]. \qquad (5.31)
\end{aligned}$$

This completes the proof.

Acknowledgement

Y.-Z.Z would like to thank Australia Research Council IREX programme for an Asia-Pacific Link Award and Institute of Modern Physics of Northwest University for hospitality.The financial support from Australian Research Council large, small and QEII fellowship grants is also gratefully acknowledged.

References

[1] C. Itzykson, H. Saleur, J.-B. Zuber, eds:*Conformal invariance and apllications to statistical mechanics*, World Scientific, Singapore, 1988.

[2] M. Jimbo, T. Miwa, *Alegbraic Analysis of Solvable Lattice Models*, CBMS Regional Conference Series in Mathematics, **Vol.85**, AMS, 1994.

[3] V.G. Drinfeld, Sov. Math. Dokl. **36** (1988) 212.

[4] H.Yamane, e-print q-alg/9603015.

[5] Y.-Z. Zhang, J. Phys. **A30** (1997) 8325; Phys. Lett. **A234** (1997) 20.

[6] J.-F. Cai, S.K. Wang, K. Wu, W.-Z. Zhao, J. Phys. **A31** (1998) 1989.

[7] M.D. Gould, Y.-Z. Zhang, Lett. Math. Phys.**44** (1998) 291.

[8] J. Ding, e-print math.QA/9905086.

[9] N.Yu. Reshetikhin, M.A. Semenov-Tian-Shansky, Lett. Math. Phys. **19** (1990) 133.

[10] J. Ding, I.B. Frenkel, Commun. Math. Phys. **155** (1993) 277.

[11] H. Awata, S. Odake, J. Shiraishi, Commun. Math. Phys. **162** (1994) 61.

[12] K. Kimura, J. Shiraishi, J. Uchiyama, Commun. Math. Phys. **188** (1997) 367.

[13] Y.-Z. Zhang, e-print math.QA/9812084.